

### Appendix SA10.3 Brief History of Leontief Inverses with Errors in the Coefficients of A

A large literature exists on “sensitivity” or “error” input-output analysis. Much of this is concerned with the situation in which all (or many) elements in  $\mathbf{A}$  are perturbed to  $a_{ij} + \Delta a_{ij}$ . This is where Woodbury’s extension (1950) of Sherman and Morrison’s results (1949, 1950) becomes relevant. Published works on how errors in an  $\mathbf{A}$  matrix influence elements in the associated Leontief inverse frequently cite results in Dwyer and Waugh (1953) and Evans (1954). An article by Henderson and Searle (1981) provides a comprehensive and unified treatment on inverses of sums of matrices. It is not concerned specifically with input-output models and does not cite either of these earlier studies. Henderson and Searle’s results, when applied to the input-output case, show that what may appear to be differing results in the input-output literature are in fact simply the same wine in different bottles.

#### A10.3.1 Mathematical Background

Much of the work on alternative expressions for the inverse of the sum of matrices is grounded in results on inverses of partitioned matrices. Recall the two results in Appendix

A for the inverse of  $\begin{bmatrix} \mathbf{E} & \mathbf{F} \\ \mathbf{G} & \mathbf{H} \end{bmatrix}$ , which we labeled  $\begin{bmatrix} \mathbf{S} & \mathbf{T} \\ \mathbf{U} & \mathbf{V} \end{bmatrix}$ ; that is,

$$\begin{bmatrix} \mathbf{E} & \mathbf{F} \\ \mathbf{G} & \mathbf{H} \end{bmatrix} \begin{bmatrix} \mathbf{S} & \mathbf{T} \\ \mathbf{U} & \mathbf{V} \end{bmatrix} = \mathbf{I} = \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{I} \end{bmatrix}$$

Building on the assumption that  $\mathbf{E}^{-1}$  is known, we have

$$\begin{aligned} \mathbf{S} &= \mathbf{E}^{-1}(\mathbf{I} - \mathbf{F}\mathbf{U}) & \mathbf{T} &= -\mathbf{E}^{-1}\mathbf{F}\mathbf{V} \\ \mathbf{U} &= -\mathbf{V}\mathbf{G}\mathbf{E}^{-1} & \mathbf{V} &= (\mathbf{H} - \mathbf{G}\mathbf{E}^{-1}\mathbf{F})^{-1} \end{aligned} \tag{A10.3.1}$$

[This is (A.5) in Appendix A.] Alternative expressions result if we begin with the assumption that  $\mathbf{H}^{-1}$  is known. This leads to

$$\begin{aligned} \mathbf{S} &= (\mathbf{E} - \mathbf{F}\mathbf{H}^{-1}\mathbf{G})^{-1} & \mathbf{T} &= -\mathbf{S}\mathbf{F}\mathbf{H}^{-1} \\ \mathbf{U} &= -\mathbf{H}^{-1}\mathbf{G}\mathbf{S} & \mathbf{V} &= \mathbf{H}^{-1}(\mathbf{I} - \mathbf{G}\mathbf{T}) \end{aligned} \tag{A10.3.2}$$

[This is (A.6).]

From the observation that corresponding expressions for  $\mathbf{S}$ ,  $\mathbf{T}$ ,  $\mathbf{U}$  or  $\mathbf{V}$  in (A10.3.1) and (A10.3.2) must be equal, we equate the two expressions for  $\mathbf{U}$ , giving

$$(\mathbf{H} - \mathbf{G}\mathbf{E}^{-1}\mathbf{F})^{-1}\mathbf{G}\mathbf{E}^{-1} = \mathbf{H}^{-1}\mathbf{G}(\mathbf{E} - \mathbf{F}\mathbf{H}^{-1}\mathbf{G})^{-1} \tag{A10.3.3}$$

Consider two special cases of this result.

(a) Let  $\mathbf{E} = \mathbf{I}$ ,  $\mathbf{F} = -\mathbf{I}$ ,  $\mathbf{G} = \boldsymbol{\alpha}$  and  $\mathbf{H} = \mathbf{I}$ , then (A10.3.3) is

$$(\mathbf{I} + \boldsymbol{\alpha})^{-1} \boldsymbol{\alpha} = \boldsymbol{\alpha}(\mathbf{I} + \boldsymbol{\alpha})^{-1} \quad (\text{A10.3.4})$$

(b) Let  $\mathbf{E} = \mathbf{I}$ ,  $\mathbf{F} = -\boldsymbol{\beta}$ ,  $\mathbf{G} = \boldsymbol{\alpha}$  and  $\mathbf{H} = \mathbf{I}$ , then (A10.3.3) is

$$(\mathbf{I} + \boldsymbol{\alpha}\boldsymbol{\beta})^{-1} \boldsymbol{\alpha} = \boldsymbol{\alpha}(\mathbf{I} + \boldsymbol{\beta}\boldsymbol{\alpha})^{-1} \quad (\text{A10.3.5})$$

Further,

$$(\mathbf{I} + \boldsymbol{\alpha})^{-1}(\mathbf{I} + \boldsymbol{\alpha}) = \mathbf{I} \Rightarrow (\mathbf{I} + \boldsymbol{\alpha})^{-1} = \mathbf{I} - (\mathbf{I} + \boldsymbol{\alpha})^{-1} \boldsymbol{\alpha} = \mathbf{I} - \boldsymbol{\alpha}(\mathbf{I} + \boldsymbol{\alpha})^{-1} \quad (\text{A10.3.6})$$

[from (A10.3.4)].

We now address the general problem of  $(\mathbf{M} + \mathbf{PNQ})^{-1}$ . Straightforward algebra leads to

$$(\mathbf{M} + \mathbf{PNQ})^{-1} = (\mathbf{M} + \mathbf{MM}^{-1}\mathbf{PNQ})^{-1} = [\mathbf{M}(\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})]^{-1} = (\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})^{-1} \mathbf{M}^{-1}$$

From (A10.3.4) and (A10.3.6), with  $\boldsymbol{\alpha} = \mathbf{M}^{-1}\mathbf{PNQ}$ ,

$$(\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})^{-1} = \mathbf{I} - (\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})^{-1} \mathbf{M}^{-1}\mathbf{PNQ}$$

and so

$$\begin{aligned} (\mathbf{M} + \mathbf{PNQ})^{-1} &= [\mathbf{I} - (\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})^{-1} \mathbf{M}^{-1}\mathbf{PNQ}] \mathbf{M}^{-1} \\ &= [\mathbf{M}^{-1} - (\mathbf{I} + \mathbf{M}^{-1}\mathbf{PNQ})^{-1} \mathbf{M}^{-1}\mathbf{PNQ} \mathbf{M}^{-1}] \end{aligned} \quad (\text{A10.3.7})$$

This is the foundation for Henderson and Searle's alternative expressions.

### *A10.3.2 Application to Leontief Inverses*

Let  $\Delta\mathbf{A}$  represent a matrix of (additive) errors (or perturbations) in the elements of a direct input coefficients matrix, so  $\mathbf{A}^* = \mathbf{A} + \Delta\mathbf{A}$ . Also,  $\mathbf{L} = (\mathbf{I} - \mathbf{A})^{-1}$  and

$$(\mathbf{I} - \mathbf{A}^*) = (\mathbf{I} - \mathbf{A}) - \Delta\mathbf{A}, \quad \mathbf{L}^* = (\mathbf{I} - \mathbf{A}^*)^{-1} = (\mathbf{I} - \mathbf{A} - \Delta\mathbf{A})^{-1}$$

Our interest is in expressing  $\mathbf{L}^*$  as a function of  $\mathbf{L}$  and  $\Delta\mathbf{A}$ <sup>1</sup>. Henderson and Searle (1981) present a number of such expressions. We particularize the results in (A10.3.7) to the Leontief inverse case by letting

$$\mathbf{M} = (\mathbf{I} - \mathbf{A}), \quad \mathbf{N} = \Delta\mathbf{A}, \quad \mathbf{P} = -\mathbf{I} \quad \text{and} \quad \mathbf{Q} = \mathbf{I} \quad [\text{and recalling that } \mathbf{L} = (\mathbf{I} - \mathbf{A})^{-1}]$$

---

<sup>1</sup> Many publications use  $\mathbf{D}$  (for "difference" matrix) or  $\mathbf{E}$  (for "error" matrix) in place of  $\Delta\mathbf{A}$ .

so that  $(\mathbf{M} + \mathbf{PNQ})^{-1} = (\mathbf{I} - \mathbf{A} - \Delta\mathbf{A})^{-1}$ . Of interest are the following from Henderson and Searle; as the authors note, they display an interesting pattern.<sup>2</sup>

$$\mathbf{L}^* = \mathbf{L} + [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1} \mathbf{L}(\Delta\mathbf{A})\mathbf{L} \quad (\text{A10.3.8})$$

This is exactly (A10.3.7), with the appropriate substitutions.

Repeatedly using (A10.3.5) generates the remaining three results of interest:

$$\mathbf{L}^* = \mathbf{L} + \mathbf{L}[\mathbf{I} - (\Delta\mathbf{A})\mathbf{L}]^{-1}(\Delta\mathbf{A})\mathbf{L} \quad (\text{A10.3.9})$$

$$\mathbf{L}^* = \mathbf{L} + \mathbf{L}(\Delta\mathbf{A})[\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1}\mathbf{L} \quad (\text{A10.3.10})$$

$$\mathbf{L}^* = \mathbf{L} + \mathbf{L}(\Delta\mathbf{A})\mathbf{L}[\mathbf{I} - (\Delta\mathbf{A})\mathbf{L}]^{-1} \quad (\text{A10.3.11})$$

The various (seemingly different) relationships presented in published input-output work are simply variations on these results.

*Dwyer and Waugh (1953)*

In Dwyer and Waugh (1953)<sup>3</sup> we find

$$\mathbf{L}^* = \mathbf{L}[\mathbf{I} - (\Delta\mathbf{A})\mathbf{L}]^{-1} \quad (\text{A10.3.12})$$

$$\mathbf{L}^* = [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1}\mathbf{L} \quad (\text{A10.3.13})$$

$$\mathbf{L}^* - \mathbf{L} = \mathbf{L}(\Delta\mathbf{A})\mathbf{L}[\mathbf{I} - (\Delta\mathbf{A})\mathbf{L}]^{-1} \quad (\text{A10.3.14})$$

$$\mathbf{L}^* - \mathbf{L} = (\mathbf{I} - \mathbf{L}(\Delta\mathbf{A}))^{-1}\mathbf{L}(\Delta\mathbf{A})\mathbf{L} \quad (\text{A10.3.15})$$

The latter two results are just (A10.3.11) and (A10.3.8), respectively. (Clearly Dwyer and Waugh's results predate the Henderson and Searle article, but they are not mentioned in it.) It is straightforward to show that (A10.3.9) and (A10.3.12) are equivalent, from (A10.3.4) with  $\mathbf{a} = -(\Delta\mathbf{A})\mathbf{L}$ . Exactly the same sort of reasoning, with  $\mathbf{a} = -\mathbf{L}(\Delta\mathbf{A})$ , shows the equivalence of (A10.3.10) and (A10.3.13).

*Evans (1954)*

The results of interest presented in Evans [1954, (1.5) and (1.6), respectively] are

$$\mathbf{L}^* = \mathbf{L}[\mathbf{I} - (\Delta\mathbf{A})\mathbf{L}]^{-1} \quad (\text{A10.3.16})$$

and

$$\mathbf{L}^* = [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1}\mathbf{L} \quad (\text{A10.3.17})$$

These are (A10.3.12) and (A10.3.13) from Dwyer and Waugh and hence equivalent to (A10.3.9) and (A10.3.10). (Evans is also not cited by Henderson and Searle.)

*West (1982)*

Here we find [West's (4)]

<sup>2</sup> These are Equations (21), (22), (24) and (26) in the article, with notation adjusted for the present input-output case.

<sup>3</sup> These are Equations (3.2), (3.2)', (3.6) and (3.6)' in the article.

$$\mathbf{L}^* = \mathbf{L} + [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1} \mathbf{L}(\Delta\mathbf{A})\mathbf{L} \quad (\text{A10.3.18})$$

which is Dwyer and Waugh's (A10.3.15) and hence (A10.3.8) from Henderson and Searle. [West cites Evans but not Dwyer and Waugh and then provides his own derivation of the result in (A10.3.18).]<sup>4</sup>

### *Sherman-Morrison and Sonis-Hewings*

The matrix statement in (A10.3.8),

$$\mathbf{L}^* = \mathbf{L} + [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1} \mathbf{L}(\Delta\mathbf{A})\mathbf{L} = \mathbf{L} + \Delta\mathbf{L}$$

is one possible generalization of the Sherman-Morrison result that we saw in Appendix 7.1 in (A7.1.4),

$$l_{rs(j)}^* = l_{rs} + \frac{l_{ri}l_{js}\Delta a_{ij}}{1 - l_{ji}\Delta a_{ij}} = l_{rs} + \Delta l_{rs(j)}$$

To demonstrate the equivalence of the general case, namely of  $\Delta l_{rs(j)}$  in  $\Delta\mathbf{L}$  [in (A10.3.8)] and its counterpart in (A7.1.4), requires fairly complicated notation, including representing elements in the inverse in (A10.3.8).

Instead, we illustrate the point for a three-sector model with a change only in  $a_{23}$  ;

that is,  $\Delta\mathbf{A} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \Delta a_{23} \\ 0 & 0 & 0 \end{bmatrix}$ . This will generate (the interested reader should check this)

$$\mathbf{L}(\Delta\mathbf{A}) = \begin{bmatrix} 0 & 0 & l_{12}\Delta a_{23} \\ 0 & 0 & l_{22}\Delta a_{23} \\ 0 & 0 & l_{32}\Delta a_{23} \end{bmatrix} \text{ and } \mathbf{L}(\Delta\mathbf{A})\mathbf{L} = \begin{bmatrix} l_{12}l_{31} & l_{12}l_{32} & l_{12}l_{33} \\ l_{22}l_{31} & l_{22}l_{32} & l_{22}l_{33} \\ l_{32}l_{31} & l_{32}l_{32} & l_{32}l_{33} \end{bmatrix} (\Delta a_{23})$$

Notice that  $\mathbf{L}(\Delta\mathbf{A})\mathbf{L} = \mathbf{F}[2,3](\Delta a_{23})$ , using the "field of influence" notation of Sonis and Hewings (section 7.3.6). In this small example,

$$[\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1} = \begin{bmatrix} 1 & 0 & l_{12}\Delta a_{23} \\ 0 & 1 & l_{22}\Delta a_{23} \\ 0 & 0 & 1 \end{bmatrix} \left( \frac{1}{[1 - l_{32}\Delta a_{23}]} \right)$$

This leads, finally, to

$$\Delta\mathbf{L}_{(23)} = [\mathbf{I} - \mathbf{L}(\Delta\mathbf{A})]^{-1} \mathbf{L}(\Delta\mathbf{A})\mathbf{L} = \begin{bmatrix} l_{12}l_{31} & l_{12}l_{32} & l_{12}l_{33} \\ l_{22}l_{31} & l_{22}l_{32} & l_{22}l_{33} \\ l_{32}l_{31} & l_{32}l_{32} & l_{32}l_{33} \end{bmatrix} \left( \frac{\Delta a_{23}}{[1 - l_{32}\Delta a_{23}]} \right) = \mathbf{F}[2,3] \left( \frac{\Delta a_{23}}{[1 - l_{32}\Delta a_{23}]} \right)$$

---

<sup>4</sup> Lahr (2001; Appendix 1) contains a very brief discussion of Dwyer and Waugh (1953), Evans (1954) and West (1982).

Clearly, any element  $\Delta l_{rs(23)}$  in this expression has exactly the structure of the Sherman-Morrison result in (A7.1.4).

We finish with a variant of the numerical example in section 7.3.4 in the text, where

we had  $\mathbf{A} = \begin{bmatrix} .15 & .25 & .05 \\ .20 & .05 & .40 \\ .30 & .25 & .05 \end{bmatrix}$  and  $\mathbf{L} = \begin{bmatrix} 1.3651 & .4253 & .2509 \\ .5273 & 1.3481 & .5954 \\ .5698 & .4890 & 1.2885 \end{bmatrix}$ . To see how the algebra

plays out, let  $a_{23}$  increase by ten percent— $\Delta a_{23} = 0.04$ —so  $\Delta \mathbf{A} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0.04 \\ 0 & 0 & 0 \end{bmatrix}$ . Applying

(A10.3.8), we need

$$\mathbf{L}(\Delta \mathbf{A}) = \begin{bmatrix} 0 & 0 & 0.0170 \\ 0 & 0 & 0.0539 \\ 0 & 0 & 0.0196 \end{bmatrix}, \quad \mathbf{L}(\Delta \mathbf{A})\mathbf{L} = \begin{bmatrix} 0.0097 & 0.0083 & 0.0219 \\ 0.0307 & 0.0264 & 0.0695 \\ 0.0111 & 0.0096 & 0.0252 \end{bmatrix}$$

$$[\mathbf{I} - \mathbf{L}(\Delta \mathbf{A})]^{-1} = \begin{bmatrix} 1 & 0 & 0.0173 \\ 0 & 1 & 0.0550 \\ 0 & 0 & 1.0200 \end{bmatrix}$$

Then

$$\Delta \mathbf{L}_{23} = [\mathbf{I} - \mathbf{L}(\Delta \mathbf{A})]^{-1} \mathbf{L}(\Delta \mathbf{A})\mathbf{L} = \begin{bmatrix} 0.0099 & 0.0085 & 0.0224 \\ 0.0313 & 0.0269 & 0.0709 \\ 0.0114 & 0.0098 & 0.0257 \end{bmatrix}$$

Alternatively, with

$$\mathbf{F}[2,3] = \begin{bmatrix} 0.2423 & 0.2080 & 0.5480 \\ 0.7682 & 0.6593 & 1.7370 \\ 0.2787 & 0.2392 & 0.6302 \end{bmatrix} \quad \text{and} \quad \Delta a_{23} / [1 - l_{32} \Delta a_{23}] = 0.0408$$

we have

$$\Delta \mathbf{L}_{23} = \mathbf{F}[2,3] \Delta a_{23} / [1 - l_{32} \Delta a_{23}] = \begin{bmatrix} 0.0099 & 0.0085 & 0.0224 \\ 0.0313 & 0.0269 & 0.0709 \\ 0.0114 & 0.0098 & 0.0257 \end{bmatrix}$$

Then

$$\mathbf{L} + \Delta \mathbf{L}_{23} = \begin{bmatrix} 1.3750 & 0.4337 & 0.2733 \\ 0.5587 & 1.3750 & 0.6662 \\ 0.5812 & 0.4988 & 1.3142 \end{bmatrix}$$

and, as expected, for  $\mathbf{A}^* = \begin{bmatrix} 0.15 & 0.25 & 0.05 \\ 0.20 & 0.05 & 0.44 \\ 0.30 & 0.25 & 0.05 \end{bmatrix}$ ,

$$\mathbf{L}^* = (\mathbf{I} - \mathbf{A}^*)^{-1} = \begin{bmatrix} 1.3750 & 0.4337 & 0.2733 \\ 0.5587 & 1.3750 & 0.6662 \\ 0.5812 & 0.4988 & 1.3142 \end{bmatrix}$$

## References

- Dwyer, Paul S. and Frederick V. Waugh. 1953. "On Errors in Matrix Inversion," *Journal of the American Statistical Association*, **48**, 289-319.
- Evans, W. Duane. 1954. "The Effect of Structural Matrix Errors on Interindustry Relations Estimates," *Econometrica*, **22**, 461-480.
- Henderson, H. V. and S. R. Searle. 1981. "On Deriving the Inverse of a Sum of Matrices," *SIAM Review*, **23**, 53-60.
- Lahr, Michael L. 2001. "A Strategy for Producing Regional Input-Output Tables," in Michael L. Lahr and Erik Dietzenbacher (eds.), *Input-Output Analysis: Frontiers and Extensions*. New York: Palgrave, pp. 211-242.
- Sherman, Jack and Winifred J. Morrison. 1949. "Adjustment of an Inverse Matrix Corresponding to Changes in the Elements of a Given Column or a Given Row of the Original Matrix (Abstract)," *Annals of Mathematical Statistics*, **20**, 621.
- \_\_\_\_\_. 1950. "Adjustment of an Inverse Matrix Corresponding to a Change in One Element of a Given Matrix," *Annals of Mathematical Statistics*, **21**, 124-127.
- Sonis, Michael and Geoffrey J. D. Hewings. 1989. "Error and Sensitivity Input-Output Analysis: A New Approach," in Ronald E. Miller, Karen R. Polenske and Adam Z. Rose (eds.), *Frontiers of Input-Output Analysis*. New York: Oxford University Press, pp. 232-244.
- West, Guy R. 1982. "Sensitivity and Key Sector Analysis in Input-Output Models," *Australian Economic Papers*, **21**, 365-378.
- Woodbury, Max A. 1950. "Inverting Modified Matrices." Memorandum Report No. 42. Statistical Research Group, Princeton University, Princeton, NJ.